

## Orthogeriatrics—Clinical Summary Document

# Geriatric fracture patient: Comprehensive geriatric assessment

Preoperative assessment of the functional, cognitive, and physical health status of elderly adults is essential to determine procedural risk, anticipate complications, and predict prognosis. Attention to these areas will assist in optimizing operative and postoperative treatment plans, as well as enhancing patient outcomes.

### Key Points:

- Specific evaluation of functional, cognitive, and physical health status is essential for all geriatric patients.
- There are simple bedside tools available to assist with these assessments (eg, Parker Mobility Scale, Mini-Cog, or Confusion Assessment Method).
- Evaluation of preoperative mobility for dementia, delirium, and polypharmacy is important for maximizing outcomes and predicting prognosis.

### Functional status/ functional ability

For most patients, functional status (along with cognitive status) is a more robust predictor of surgical outcomes than physical health and comorbidity. Functional status can be rapidly assessed with a number of validated approaches including the following:

#### 1. ADL/IADL assessment (for all patients)

Activities of daily living (ADLs) and instrumental activities of daily living (IADLs) are simple ways to assess a patient's recent functional status. ADL and IADL deficiencies predict poorer prognoses.

#### ADLs

Bathing

Dressing

Toileting/continence

Transferring (bed to chair)

Grooming

Feeding

#### IADLs

Self-administration of medications

Food shopping

Meal preparation

Managing finances

Using telephone

Driving

Housekeeping

Laundry

2. **New Mobility Scale (Parker Mobility Scale)**

This is a simple evaluation of prefracture mobility and has been shown to correlate with postoperative outcomes.

MOBILITY	No Difficulty	With Aid	With help from another person	Not at all
Able to get about the house	3	2	1	0
Able to get out of the house	3	2	1	0
Able to go shopping	3	2	1	0
One year mortality for NMS:		2	1	0

3. **Timed Up and Go Test (TUG)** (for ambulatory patients, but can also be used postoperatively).

This simple functional test measures the speed at which the patient can arise from a standard armchair, walk 3 meters, turn around, and sit back down in the chair. Patients may use any customary assistive device. After performing one practice trial, the time results of two subsequent trials are averaged. Time > 14 seconds at baseline is considered a high risk for falls. TUG time > 56 seconds 4 days after hemiarthroplasty predicts dependence on assistive devices at 2 years.

4. **Additional relevant questions to screen for functional impairment:**

Have you had any weight loss in the past 6 months?

Have you had any falls in the past 6 months?

## Cognitive status/cognitive ability

1. Screen for dementia in patients with stable cognition using the **Mini-Cognitive Assessment Instrument (Mini-Cog)**. The Mini Cog enables rapid screening for dementia in a patient at their cognitive baseline (not delirious). This simple assessment has been well validated and compared to more comprehensive assessments such as the Mini-Mental Status examination.

**Step 1.** Ask the patient to repeat three unrelated words

**Step 2.** Ask the patient to draw a clock, and set the time to 10 minutes after 11 (11:10). A correct response is to draw a circle with the numbers in their approximately correct positions, with the hands pointing to the 11 and 2. (two points for correct clock, zero point for incorrect clock)

**Step 3.** Ask the patient to recall the three items from **Step 1**. (one point for each item recalled correctly)

**Scoring:**

**Zero to two points:** Positive screen for dementia

**Three to five points:** Negative screen for dementia

2. Evaluate delirium (in patients with unstable/changing cognition) with the Confusion Assessment Method (CAM). Delirium can have reversible causes (eg, pain, hypoxia, medications, infection, constipation, or urinary retention) and is also a marker for poor physical and cognitive outcomes. It is more common in patients with an underlying baseline cognitive impairment. The CAM enables rapid diagnosis of delirium in a patient with acutely altered mental status. The information for this assessment can typically be obtained during the course of a normal history and consists of four criteria related to the altered mental status:

1. **Acute onset/fluctuating course** (chronic cognitive impairment is not delirium)
2. **Inattention** (difficulty keeping track of current conversation)
3. **Disorganized thinking** (rambling conversation, incoherent ideas, hallucinations, or delusions)
4. **Altered level of consciousness** (hyperalert or lethargic)

Delirium is diagnosed when criteria **1 and 2, and** either **3 or 4** are present.

## Physical health and assessment for comorbidity

The history and physical exam continue to be the cornerstone for optimizing the geriatric fracture patient for surgery. Rapidly identifying unstable medical conditions is essential, and stopping some medications that are problematic in this acute setting (eg, anticholinergic medications such as diphenhydramine or many antihypertensive medications) is important to perioperative and postoperative outcomes.

Focused questions and physical exam with attention to the following:

- Circumstances of fall and injury (to identify medical contributions to fall)
- Screening for other injuries
- Past medical history and recent hospitalizations
- Previous screening, diagnosis, and treatment for osteoporosis
- Previous surgical, anesthetic complications, including delirium
- Medication review and possible effects of polypharmacy
- Physical exam to exclude active medical problems (eg, pulmonary edema, stroke, aortic stenosis, or hemodynamic instability)
- Basic laboratory assessment (blood count, basic blood chemistry analysis, electrocardiogram, or 25-OH vitamin D levels)
- For anemic patients without symptoms, keeping postoperative hemoglobin > 8 g/dL appears to produce similar outcomes to more liberal transfusion strategies.

## Recommended readings

1. **Malani PN.** Functional status assessment in the preoperative evaluation of older adults. *JAMA*. 2009 Oct 14;302(14):1582–1583.
2. **Svensson O, Strömberg L, Ohlén G, et al.** Prediction of the outcome after hip fracture in elderly patients. *J Bone Joint Surg Br*. 1996 Jan;78(1):115–118.
3. **Parker MJ, Palmer CR.** A new mobility score for predicting mortality after hip fracture. *J Bone Joint Surg Br*. 1993 Sep; 75(5):797–798.
4. **Laflamme GY, Rouleau DM, Leduc S, et al.** The Timed Up and Go test is an early predictor of functional outcome after hemiarthroplasty for femoral neck fracture. *J Bone Joint Surg Am*. 2012 Jul 3;94(13):1175–1179.
5. **Borson S, Scanlan J, Brush M, et al.** The mini-cog: a cognitive 'vital signs' measure for dementia screening in multi-lingual elderly. *Int J Geriatr Psychiatry*. 2000 Nov;15(11):1021–1027.

6. **Inouye SK, van Dyck CH, Alessi CA, et al.** Clarifying confusion: the confusion assessment method. A new method for detection of delirium. *Ann Intern Med.* 1990 Dec 15;113(12):941–948.
  7. **Carson JL, Terrin ML, Noveck H, et al.** Liberal or restrictive transfusion in high-risk patients after hip surgery. *N Engl J Med.* 2011 Dec 29;365(26):2453–2462.
- 

## Disclaimer

Production: AO Foundation, Switzerland

### Hazards

Great care has been taken to maintain the accuracy of the information contained in this publication. However, the publisher, and/or the distributor, and/or the editors, and/or the authors cannot be held responsible for errors or any consequences arising from the use of the information contained in this publication. Contributions published under the name of individual authors are statements and opinions solely of said authors and not of the publisher, and/or the distributor, and/or the AO Group.

The products, procedures, and therapies described in this work are hazardous and are therefore only to be applied by certified and trained medical professionals in environments specially designed for such procedures. No suggested test or procedure should be carried out unless, in the user's professional judgment, its risk is justified.

Whoever applies products, procedures, and therapies shown or described in this work will do this at their own risk. Because of rapid advances in the medical sciences, AO recommends that independent verification of diagnosis, therapies, drugs, dosages, and operation methods should be made before any action is taken.

Although all advertising material which may be inserted into the work is expected to conform to ethical (medical) standards, inclusion in this publication does not constitute a guarantee or endorsement by the publisher regarding quality or value of such product or of the claims made of it by its manufacturer.

### Legal restrictions

This work was produced by AO Foundation, Switzerland. All rights reserved. This publication, including all parts thereof, is legally protected by copyright. Any use, exploitation or commercialization outside the narrow limits set forth by copyright legislation and the restrictions on use laid out below, without the publisher's consent, is illegal and liable to prosecution. This applies in particular to photostat reproduction, copying, scanning or duplication of any kind, translation, preparation of microfilms, electronic data processing, and storage such as making this publication available on Intranet or Internet. Some of the products, names, instruments, treatments, logos, designs, etc. referred to in this publication are also protected by patents and trademarks or by other intellectual property protection laws (e.g. "AO", "ASIF", "AO/ASIF", TRIANGLE/GLOBE Logo" are registered trademarks) even though specific reference to this fact is not always made in the text. Therefore, the appearance of a name, instrument, etc. without designation as proprietary is not to be construed as a representation by the publisher that it is in the public domain.

Restrictions on use: The rightful owner of an authorized copy of this work may use it for educational and research purposes only. Single images or illustrations may be copied for research or educational purposes only. The images or illustrations may not be altered in any way and need to carry the following statement of origin "Copyright by AO Foundation, Switzerland".

Copyright © 2014 by AO Foundation, Switzerland, Clavadelstrasse 8, CH-7270 Davos Platz